

UNCLASSIFIED

AD 298 258

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

298 258

Technical Report

ON CONVERGENCE PROOFS FOR PERCEPTRONS

Prepared for:

OFFICE OF NAVAL RESEARCH
WASHINGTON, D.C.


CONTRACT Nonr 3438(00)

By: Albert B. J. Novikoff

CATALOGED BY
AS AD NO. 1

STANFORD RESEARCH INSTITUTE

MENLO PARK, CALIFORNIA





January 1963

Technical Report

ON CONVERGENCE PROOFS FOR PERCEPTRONS

Prepared for:

OFFICE OF NAVAL RESEARCH
WASHINGTON, D.C.

CONTRACT Nonr 3438(00)

By: Albert B. J. Novikoff

SRI Project No. 3605

Approved:

C. A. ROSEN, MANAGER APPLIED PHYSICS LABORATORY

J. D. NOE, DIRECTOR ENGINEERING SCIENCES DIVISION

Copy No. 3

ABSTRACT

A short proof is given of the convergence (in a finite number of steps) of an algorithm for adjusting weights in a single-threshold device. The algorithm in question can be interpreted as the error-correction procedure introduced by Rosenblatt for his " α -Perceptron." The proof presented extends the basic idea to continuous as well as discrete cases, and is interpreted geometrically.

CONTENTS

ABSTRACT	ii
I INTRODUCTION	1
II STATEMENT OF THE THEOREM	2
III PROOF OF THEOREM	4
IV AN ANALOGOUS THEOREM	6
V GEOMETRICAL INTERPRETATION	8
ACKNOWLEDGMENT	11
REFERENCES	12

I INTRODUCTION

The purpose of this report is to exhibit an extremely short and, more notably, transparent proof of a theorem concerning perceptrons. The theorem itself must now be considered one of the most basic theorems about perceptrons, and indeed, is among the first theorems proved by Rosenblatt and his collaborators. It also enjoys the peculiar distinction of being one of the most often re-proved results in the field. The succession of proofs now available progresses from somewhat cloudy statements (which at one time caused doubt among "reasonable men" that the theorem was true) to comparatively crisp statements of a purely mathematical nature which nonetheless use more print than is strictly necessary.¹⁶

More to the point, latter-day proofs fail to enunciate a simple principle involved. This principle permits one to modify the hypotheses in a variety of ways and secure similar results; it may well be useful in establishing genuinely new theorems of like character. We therefore present our proof in its entirety, in part to verify our claim that it is as short a line as can be drawn from hypotheses to conclusion, and also with the hope of terminating an already lengthy process of successive refinements. In addition, we prove a related theorem which is the continuous analogue of the perceptron theorem, and we indicate that various other theorems may be obtained by appropriately modifying the hypotheses. We also discuss a geometrical interpretation of the perceptron theorem in terms of a convex cone and its dual.

II STATEMENT OF THE THEOREM

Whereas previous proofs of the theorem appealed to a structure, called by Rosenblatt and his co-workers an α -perceptron, the present theorem, proof, and discussion apply without modification to a structure consisting of a single threshold element acting on a weighted set of inputs.

Theorem: Let w_1, \dots, w_N be a set of vectors in a Euclidean space of fixed finite dimension, satisfying the hypothesis that there exists a vector y such that

$$(w_i, y) > \theta > 0 \quad i = 1, \dots, N \quad (1)$$

Consider the infinite sequence $w_{i_1}, w_{i_2}, w_{i_3}, \dots$, $1 \leq i_k \leq N$ for every k , such that each vector w_1, \dots, w_N occurs infinitely often. Recursively construct a sequence of vectors $v_0, v_1, \dots, v_n, \dots$ as follows:

v_0 is arbitrary

$$v_n = \begin{cases} v_{n-1} & \text{if } (w_{i_n}, v_{n-1}) > \theta \\ v_{n-1} + w_{i_n} & \text{if } (w_{i_n}, v_{n-1}) \leq \theta \end{cases} \quad (2)$$

The sequence $\{v_n\}$ is convergent—i.e., for some index m , $v_m = v_{m+1} = v_{m+2} = \dots = \tilde{v}$.

Remarks:

(1) In particular, the theorem insures that $(w_i, \tilde{v}) > \theta$ for $i = 1, \dots, N$, since each w_i occurs arbitrarily far out in the sequence $\{w_{i_k}\}$. It is only to obtain this consequence that we impose the restriction that each w_i occurs infinitely often in the training sequence.

(2) Theoretically, we may take $\theta = 0$ without loss of generality.¹ However, this often has the effect of smuggling in numbers of large magnitude. For this reason, we retain the general θ , but in the concluding section we do consider the relation between the general case and the case $\theta = 0$.

(3) In the private language of perceptron workers, the theorem reads as follows: A set of incoming signals is divided into two adjacent classes. A "satisfactory" assignment of weights from the associator units is defined as an assignment resulting in a response +1 for signals of Class I, and -1 for signals of Class II. The theorem asserts that no matter what assignment of weights we begin with, the process of recursively readjusting the weights by the method known as "error correction" will terminate after a finite number of corrections in a satisfactory assignment, provided such a satisfactory assignment exists. More briefly, a finite number of corrections will teach the perceptron to perform any given dichotomy of signals, if the dichotomy is within the capacity of the perceptron at all.

The definition of α -perceptron and the precise correspondence between the theorem's original verbal description and the purely mathematical assertion of the above theorem are provided by Block.¹ A brief glossary indicating the correspondence follows: the vector w_i represents the activity of the associators, including class information, when stimulus S_i is presented. The vector y , which we assume to exist, represents a "satisfactory" assignment of associator weights; y has as many components as there are associators. The sequence $\{w_{i,n}\}$ represents the "training sequence," and the rule for defining $\{v_n\}$ describes the error-correction procedure. The positive number θ is a threshold which must be exceeded for the response of the perceptron to be correct; a vector v such that $(w_i, v) > \theta$ is an assignment of associator outputs which successfully classifies the i th signal. [If $(w_i, v) \leq \theta$, then either the i th signal has been classified as belonging to the incorrect class or the perceptron has refrained from commitment, depending on whether or not the inequality is strict.]

(4) It is clear that because of Eq. (2), as n varies, the sequence v_n changes, if at all, only by the addition of one or another of the set w_1, \dots, w_N . For this reason "convergence" implies "convergence in a finite number of steps." The word "stabilizes" has been suggested to describe this kind of convergence.

III PROOF OF THEOREM

We may omit from the training sequence all terms w_{i_n} for which $v_n = v_{n-1}$, as these w_{i_n} are clearly inessential. The new training sequence is such that correction takes place at every step. Adjusting our notation, we may assume that

$$v_n = v_{n-1} + w_{i_n} \quad \text{and} \quad (w_{i_n}, v_{n-1}) \leq \theta \quad \text{for each } n. \quad (3)$$

We observe that n is the number of corrections made up to the n th step. The assertion of the theorem after this change of notation is that n can range only through a finite set of integers—that is, conditions (1) and (3) cannot continue to hold simultaneously for all $n = 1, 2, 3, \dots$.

First we show that inequality (1) alone implies the inequality

$$\|v_n\|^2 > Cn^2 \quad (4)$$

for suitable choice of the positive constant C , and n sufficiently large. Since $v_n = v_0 + w_{i_1} + \dots + w_{i_n}$, inequality (1) implies that v_n satisfies $(v_n, y) > (v_0, y) + n\theta$. Using the Cauchy-Schwartz inequality,

$$\|v_n\|^2 \geq \frac{(v_n, y)^2}{\|y\|^2} > \frac{[(v_0, y) + n\theta]^2}{\|y\|^2} = \frac{\theta^2}{\|y\|^2} \left[n + \frac{(v_0, y)}{\theta} \right]^2$$

If $(v_0, y) \geq 0$, we may choose $C = \theta^2/\|y\|^2$, and inequality (4) is satisfied for all n . If $(v_0, y) < 0$, we may choose $C = (1/4)(\theta^2/\|y\|^2)$, and inequality (4) is satisfied for $n > -2[(v_0, y)/\theta]$.

On the other hand, we show that inequalities (3) alone imply the inequality

$$\|v_n\|^2 \leq \|v_0\|^2 + (2\theta + M)n \quad (5)$$

where

$$M = \max_{i=1, \dots, N} \|w_{i_i}\|^2$$

Using inequality (3), the integer-argument function $\|v_k\|^2$ satisfies for each k the difference inequality

$$\|v_k\|^2 - \|v_{k-1}\|^2 = 2(v_{k-1}, w_{i_k}) + \|w_{i_k}\|^2 \leq 2\theta + M.$$

Adding the inequalities for $k = 1, 2, \dots, n$, we obtain inequality (5).

Clearly, inequalities (4) and (5) are incompatible for n sufficiently large.

IV AN ANALOGOUS THEOREM

The theorem of Section II is the discrete analogue of the following theorem, which may seem more intuitive: Let $v(t)$ be a curve in Euclidean n -space described by a smooth vector function of the continuous variable t , such that there exists a vector y such that

$$\left(\frac{dv}{dt}, y \right) > C > 0 \quad (1)'$$

and

$$\frac{1}{2} \frac{d}{dt} \|v(t)\|^2 = \left[\frac{dv}{dt}, v(t) \right] \leq \theta, \quad 0 \leq t < b \quad (3)'$$

There exists an upper bound for b ; in particular, inequalities (1)' and (3)' are compatible only over a finite domain on the t -axis.

The proof is virtually identical with that of the discrete case. Integrating inequality (1)' from 0 to t , we obtain

$$[v(t), y] > [v(0), y] + Ct \quad (6)$$

Integrating inequality (3)' from 0 to t , we obtain

$$\|v(t)\|^2 \leq 2\theta t + \|v(0)\|^2 \quad (7)$$

Using the Cauchy-Schwartz inequality and inequality (6), we obtain

$$\|v(t)\|^2 \geq \frac{[v(t), y]^2}{\|y\|^2} > \frac{\{[v(0), y] + Ct\}^2}{\|y\|^2} \quad (8)$$

Inequalities (7) and (8) together show that t cannot exceed the larger root of the quadratic

$$\{[v(0), y] + Ct\}^2 = \|y\|^2 \{2\theta t + \|v(0)\|^2\}$$

Remarks:

(1) Inequality (1)' means that the tangent vector to the curve lies on one side of a hyperplane. Inequality (3)' means that the rate of growth of $\|v\|^2$ is bounded.

(2) We may compare the above argument for the continuous case with the extremely familiar phenomenon that

$$\left[v(t), \frac{dv}{dt} \right] = 0 \quad (9)$$

implies

$$\|v(t)\|^2 \text{ is constant } ; \quad (10)$$

i.e., a curve whose tangent vector is always perpendicular to its position vector is constrained to lie on a sphere. Replacing the orthogonality condition (9) with the inequality (3)' results in an inequality (7) on the rate of growth of the function $f(t) = \|v(t)\|^2$, which is clearly a weakening of the condition (10) that $f(t)$ be constant.

(3) The principle involved in the theorem is the following: The condition of (3)', that the tangent vector have bounded scalar product with the position vector, clearly results in an upper bound for the instantaneous position of the curve as a function of time. On the other hand, if the tangent vectors dv/dt to the curve remain sufficiently large and do not depart too badly from colinearity, as prescribed, for example, by (1)', then a lower bound on the cumulative growth results, as in (8). This is intuitively clear: If dv/dt does not get too small, the total arclength will increase with at least a certain rate. If, on the other hand, the dv/dt are sufficiently "nearly colinear" then the serpentine path swept out by $v(t)$ cannot reverse its direction enough to prevent its over-all migration away from its starting point. The opposition of these two influences implies the termination of one of the two relations (1)' or (3)'.

We will not dwell upon the matter of how assorted variations of this theme will continue to produce assertions that t , or, in the discrete case, n , must remain bounded. Whether each of these deserves to be dignified with the name *theorem* is a moot point.

V GEOMETRICAL INTERPRETATION

We conclude with a few words about the geometrical interpretation of the assumption (1). We assume familiarity with the theory of convex sets in Euclidean vector spaces. For the most part, we state these remarks without proof. For a general introduction to this theory, see Blackwell and Girshick,⁹ and Gale.¹⁰

The *polyhedral cone*, C , with generators, w_1, \dots, w_N , is defined as all vectors of the form $\lambda_1 w_1 + \dots + \lambda_N w_N$, where $\lambda_i \geq 0$, $i = 1, 2, \dots, N$. The cone C is called *proper* if for all $v \neq 0$, C never contains both v and $-v$; or equivalently, if C , apart from its vertex, lies in the interior of a half space. The condition that there exists a vector y satisfying (1) is precisely equivalent to the condition that C is a proper cone.

We remark that requiring the existence of a vector y which satisfies (1) with $\theta > 0$, is neither stronger nor weaker than requiring the existence of a vector \tilde{y} which satisfies (1) with $\theta = 0$ —i.e., which satisfies

$$(w_i, \tilde{y}) > 0, \quad i = 1, \dots, N. \quad (1)''$$

Indeed, y itself can serve for \tilde{y} ; conversely, given θ and \tilde{y} satisfying (1)'', any sufficiently large positive multiple $y = \lambda \tilde{y}$ of \tilde{y} with

$$\lambda > \frac{\theta}{\min_{i=1, \dots, N} (w_i, \tilde{y})},$$

will satisfy (1). Condition (1)'' is the customary way of specifying that the cone C be proper.

For the continuous analogue, requiring that infinitely many vectors dv/dt satisfy (1)' with $c > 0$ is actually stronger than requiring only that dv/dt satisfy

$$\left(\frac{dv}{dt}, y \right) > 0, \quad 0 \leq t < b. \quad (1)'''$$

In fact, the left-hand side, though positive for each t , need not be bounded away from zero. If we assume $(1)'''$ to hold for $0 \leq t \leq b$ (equality permitted at b) and dv/dt to be continuous, then, as in the discrete case, it is true that $(1)'''$ and $(1)'$ are precisely equivalent.

The cone C^* of all vectors v such that $(w, v) \geq 0$ for all w in C , or equivalently such that

$$(w_i, v) \geq 0, \quad i = 1, \dots, N, \quad (10)$$

where C is the polyhedral cone generated by w_1, \dots, w_n , is called the *dual cone* of C . Its interior consists of all v for which every inequality in (10) is strict. The bigger C is the smaller C^* is, and vice versa; for example, when C is a half-space, C^* is a half-line. In general, for $n > 2$ neither need include the other. On the other hand, it is not possible to weakly separate C and C^* —i.e., there is no $z \neq 0$ such that both $(w, z) \geq 0$ for all w in C , and $(v, z) \leq 0$ for all v in C^* . If such a z did exist, then by the first inequality, z is in C^* ; then choosing v to be z in the second inequality implies that $(z, z) \leq 0$ —i.e., that $z = 0$, contrary to the assumption that $z \neq 0$.

When C is proper, C^* has an interior; indeed, the \tilde{y} of $(1)''$ is in the interior of C^* . If C^* has an interior, C and the interior of C^* overlap; if not, then a consequence would be that C and C^* could be weakly separated by a classic result on the separation of convex sets. This we have seen to be impossible.

As previously observed, if y is in the interior of C^* , then a suitable positive multiple of y will satisfy (1). Let the set of vectors satisfying (1) be denoted by D . D is a subset of the interior of C^* . The relation between (1) and the dual cone defined by (10) may be summarized as follows: (1) has a solution (i.e., D is non-empty) if and only if C^* has an interior.

The error-correction procedure is a *recursive construction* of a vector in D of the form

$$k_1 w_1 + \dots + k_N w_N$$

where k_i is the number of times w_i occurred in the irredundant training

sequence before the termination of the correction process. The mere existence of such a vector (without a construction algorithm) is assured by the fact that C and D overlap, which is a consequence of the above-mentioned overlap between C and the interior of C^* .

In general, if condition (1) is fulfilled, so that w_1, \dots, w_N generate a proper cone, it will happen that some subset of the w 's will generate the same cone (which then has the same dual C^*), and it suffices to restrict attention to this subset in constructing the training sequence. To accelerate the termination of the correction process one should use for correction those w 's which themselves are nearest the interior of C^* , and which are as long as possible. If, for example, $w_3 = w_1 + w_2$, the single addition of w_3 will accomplish as much as the successive additions of w_1 and w_2 . The question of whether a dichotomy is within the combinatorial capacity of an α -type perceptron reduces to whether or not C^* has an interior, or equivalently, whether or not C is proper. This question is discussed in a somewhat different context by Joseph and Hay,⁷ and Keller.⁸

After the work was completed it was pointed out that S. Agmon had previously considered¹¹ a variety of similar correction procedures (none, however, identical with the above) and showed their convergence in general to be at a geometric rate.

ACKNOWLEDGMENT

This report is substantially the same as a talk given at the Brooklyn Polytechnic Institute Symposium on Automata, in April 1962. In subsequent preparation of this paper as a report, several statements were rendered more explicit. Much gratitude is owed to Donna Kaylor for useful suggestions and patient cooperation, which resulted in greater clarity.

REFERENCES

- [1] H. D. Block, "The Perceptron: A Model for Brain Functioning. I," *Reviews of Modern Physics*, Vol. 34, No. 1 (January 1962).
- [2] F. Rosenblatt, Report VG-1196-G-4, Cornell Aeronautical Laboratory (February 1960).
- [3] R. D. Joseph, Tech. Memo 12, Project PARA, Cornell Aeronautical Laboratory (May 1960).
- [4] R. D. Joseph, Tech. Memo 13, Project PARA, Cornell Aeronautical Laboratory (July 1960).
- [5] R. A. Stafford, "Learning by Threshold Elements in Cascade," Tech. Note 20081, Aeronutronics (8 March 1961).
- [6] R. C. Singleton, "A Test for Linear Separability as Applied to Self-Organizing Machines," Conference on Self-Organizing Systems, May 22-24, 1962, Chicago, Illinois (Proceedings in press, Spartan Books).
- [7] T. F. Joseph and L. Hay, Tech. Memo 8, Project PARA, Cornell Aeronautical Laboratory (1960).
- [8] H. B. Keller, "Finite Automata, Pattern Recognition and Perceptron," *J. Association for Computing Machinery*, Vol. 8, No. 1 (January 1961).
- [9] D. Blackwell and M. A. Girshick, *Theory of Games and Statistical Decisions*, Sec. 2.2, Theorem 2.2.1, Corollary 2, pp. 33-36 (John Wiley & Sons, Inc., New York, N.Y., 1954).
- [10] David Gale, *The Theory of Linear Economic Models*, Chapter II (McGraw-Hill, New York, N.Y., 1960).
- [11] Samuel Agmon, "The Relaxation Method for Linear Inequalities," *Canadian J. of Math.*, Vol. 6, No. 3 (1954).

**STANFORD
RESEARCH
INSTITUTE**

**MENLO PARK
CALIFORNIA**

Regional Offices and Laboratories

Southern California Laboratories
820 Mission Street
South Pasadena, California

Washington Office
808 17th Street, N.W.
Washington 5, D.C.

New York Office
270 Park Avenue, Room 1770
New York 17, New York

Detroit Office
The Stevens Building
1025 East Maple Road
Birmingham, Michigan

European Office
Pelikanstrasse 37
Zurich 1, Switzerland

Japan Office
911 Iino Building
22, 2-chome, Uchisaiwai-cho, Chiyoda-ku
Tokyo, Japan

Representatives

Honolulu, Hawaii
Finance Factors Building
195 South King Street
Honolulu, Hawaii

London, England
19 Upper Brook Street
London, W. 1, England

Milan, Italy
Via Macedonio Melloni 40
Milano, Italy

London, Ontario, Canada
P.O. Box 782
London, Ontario, Canada

DISTRIBUTION LIST

Organization	No. of Copies
Assistant Sec. of Def. for Res. and Eng. Information Office Library Branch Pentagon Building Washington 25, D.C.	2
Armed Services Technical Information Agency Arlington Hall Station Arlington 12, Virginia	10
Chief of Naval Research Department of the Navy Washington 25, D.C. Attn: Code 437, Information Systems Branch	2
Chief of Naval Operations OP-07T-12 Navy Department Washington 25, D.C.	1
Director, Naval Research Laboratory Technical Information Officer, Code 2000 Washington 25, D.C.	6
Commanding Officer, Office of Naval Research Navy #100, Fleet Post Office New York, New York	10
Commanding Officer, ONR Branch Office 346 Broadway New York 13, New York	1
Commanding Officer, ONR Branch Office 495 Summer Street Boston 10, Massachusetts	1
Bureau of Ships Department of the Navy Washington 25, D.C. Attn: Code 607A NTDS	1
Bureau of Naval Weapons Department of the Navy Washington 25, D.C. Attn: RAAV Avionics Division	1

<u>Organization</u>	<u>No. of Copies</u>
Bureau of Naval Weapons Department of the Navy Washington 25, D.C. Attn: RMWC Missile Weapons Control Div.	1
Bureau of Ships Department of the Navy Washington 25, D.C. Attn: Communications Branch Code 686	1
Naval Ordnance Laboratory White Oaks Silver Spring 19, Maryland Attn: Technical Library	1
David Taylor Model Basin Washington 7, D.C. Attn: Technical Library	1
Naval Electronics Laboratory San Diego 52, California Attn: Technical Library	1
University of Illinois Control Systems Laboratory Urbana, Illinois Attn: D. Alpert	1
Air Force Cambridge Research Laboratories Laurence G. Hanscom Field Bedford, Massachusetts Attn: Research Library, CRX2-R	1
Technical Information Officer U.S. Army Signal Research & Dev. Lab. Fort Monmouth, New Jersey Attn: Data Equipment Branch	1
National Security Agency Fort George G. Meade, Maryland Attn: R-4, Howard Campaigne	1
U.S. Naval Weapons Laboratory Dahlgren, Virginia Attn: Head, Computation Div., G.H. Gleissner	1

<u>Organization</u>	<u>No. of Copies</u>
National Bureau of Standards Data Processing Systems Division Room 239, Bldg. 10 Washington 25, D.C. Attn: A. K. Smilow	1
Aberdeen Proving Ground, BRL Aberdeen Proving Ground, Maryland Attn: J. H. Giese, Chief Computation Lab.	1
Office of Naval Research Resident Representative Stanford University Electronics Res. Laboratory Stanford, California	1
Commanding Officer ONR, Branch Office John Crerar Library Bldg. 86 East Randolph Street Chicago 1, Illinois	1
Commanding Officer ONR Branch Office 1030 E. Green Street Pasadena, California	1
Commanding Officer ONR Branch Office 1000 Geary Street San Francisco 9, California	1
Syracuse University Electrical Eng. Dept. Syracuse 10, New York Attn: Dr. Stanford Goldman	1
Cornell University Cognitive Systems Research Program Hollister Hall Ithaca, New York Attn: Dr. Frank Rosenblatt	1
Communications Sciences Lab. University of Michigan 180 Frieze Building Ann Arbor, Michigan Attn: Gordon E. Peterson	1

<u>Organization</u>	<u>No. of Copies</u>
University of Michigan Ann Arbor, Michigan Attn: Dept. of Psychology, Prof. Arthur Melton	1
University of Michigan Ann Arbor, Michigan Attn: Dept. of Philosophy, Prof. A. W. Burks	1
University of Pennsylvania Philadelphia 4, Pennsylvania Attn: Mr. R. Duncan Luce Dept. of Psychology	1
Carnegie Institute of Technology Dept. of Psychology Pittsburgh 13, Pennsylvania Attn: Prof. Bert F. Green, Jr.	1
Massachusetts Institute of Technology Research Lab. for Electronics Cambridge, Massachusetts Attn: Dr. Marvin Minsky	1
Stanford University Stanford, California Attn: Electronics Lab., Prof. John G. Linvill	1
University of Illinois Urbana, Illinois Attn: Electrical Engrg. Dept. Prof. H. Von Foerster	1
University of California Institute of Eng. Research Berkeley 4, California Attn: Prof. A. J. Thomasian	1
University of California - LA Los Angeles 24, California Attn: Dept. of Engineering Prof. Gerald Estrin	1
Massachusetts Institute of Technology Research Laboratory of Electronics Attn: Prof. W. McCulloch	1

<u>Organization</u>	<u>No. of Copies</u>
University of Michigan Ann Arbor, Michigan Attn: Dept. of Psychology, Prof. Arthur Melton	1
University of Michigan Ann Arbor, Michigan Attn: Dept. of Philosophy, Prof. A. W. Burks	1
University of Pennsylvania Philadelphia 4, Pennsylvania Attn: Mr. R. Duncan Luce Dept. of Psychology	1
Carnegie Institute of Technology Dept. of Psychology Pittsburgh 13, Pennsylvania Attn: Prof. Bert F. Green, Jr.	1
Massachusetts Institute of Technology Research Lab. for Electronics Cambridge, Massachusetts Attn: Dr. Marvin Minsky	1
Stanford University Stanford, California Attn: Electronics Lab., Prof. John G. Linvill	1
University of Illinois Urbana, Illinois Attn: Electrical Engrg. Dept. Prof. H. Von Foerster	1
University of California Institute of Eng. Research Berkeley 4, California Attn: Prof. A. J. Thomasian	1
University of California - LA Los Angeles 24, California Attn: Dept. of Engineering Prof. Gerald Estrin	1
Massachusetts Institute of Technology Research Laboratory of Electronics Attn: Prof. W. McCulloch	1

<u>Organization</u>	<u>No. of Copies</u>
University of Illinois Urbana, Illinois Attn: John R. Pasta	1
Naval Research Laboratory Washington 25, D.C. Attn: Security Systems Code 5266, Mr. G. Abraham	1
Zator Company 140 1/2 Mt. Auburn Cambridge 38, Massachusetts Attn: R. J. Solomonoff	1
NASA Goddard Space Flight Center Washington 25, D.C. Attn: Arthur Shapiro	1
National Physical Laboratory Teddington, Middlesex England Attn: Dr. A. M. Uttley, Superintendent Autonomics Division	1
Diamond Ordnance Fuze Laboratory Connecticut Avenue & Van Ness St. Washington 25, D.C. Attn: ORDTL-012, E. W. Channel	1
Harvard University Cambridge, Massachusetts Attn: School of Applied Science Dean Harvey Brook	1
Wright Air Development Division Electronic Technology Laboratory Wright-Patterson AFB, Ohio Attn: Lt. Col. L. M. Butsch, Jr., ASRUEB	1
Laboratory for Electronics, Inc. 1079 Commonwealth Avenue Boston 15, Massachusetts Attn: Dr. H. Fuller	1

<u>Organization</u>	<u>No. of Copies</u>
Stanford Research Institute Computer Laboratory Menlo Park, California Attn: H. D. Crane	1
General Electric Co. Schnectady 5, N.Y. Attn: Library, L.M.E. Dept., Bldg. 28-501	1
The Rand Corp. 1700 Main St. Santa Monica, California Attn: Numerical Analysis Dept. Willis H. Ware	1
Carnegie Institute of Technology Pittsburgh, Pennsylvania Attn: Director, Computation Center Alan J. Perlis	1
Rome Air Development Center, RCOR DCS/Operations, USAF Griffiss Air Force Base, New York Attn: Irving J. Gabelman	1
Air Force Office of Scientific Research Directorate of Information Sciences Washington 25, D.C. Attn: Dr. Harold Wooster	1
Stanford Research Institute Menlo Park, California Attn: Dr. Charles Rosen Applied Physics Laboratory	1
L. G. Hanscom Field, AF-CRL-CRRB Bedford, Massachusetts Attn: Dr. H. H. Zschirnt	1
Department of the Army Office of the Chief of Research & Development Pentagon, Room 3D442 Washington 25, D.C. Attn: Mr. L. H. Geiger	1

<u>Organization</u>	<u>No. of Copies</u>
Bell Telephone Laboratories Murray Hill Laboratory Murray Hill, New Jersey Attn: Dr. Edward F. Moore	1
Carnegie Institute of Technology Systems and Communication Sciences Pittsburgh 13, Pennsylvania Attn: Dr. Allen Newell	1
Department of Commerce U.S. Patent Office Washington 25, D.C. Attn: Mr. Herbert R. Koller	1
Northwestern University Computer Sciences Lab. The Technological Inst. Evanston, Illinois Attn: Dr. Julius T. Tou	1
University of Pennsylvania Moore School of Electrical Engineering 200 South 33rd Street Philadelphia 4, Pennsylvania Attn: Miss Anna Louise Campion	1
University of Pennsylvania Mechanical Languages Projects Moore School of Electrical Engineering Philadelphia 4, Pennsylvania Attn: Dr. Saul Gorn, Director	1
Applied Physics Laboratory Johns Hopkins University 8621 Georgia Avenue Silver Spring, Maryland Attn: Document Library	1
Bureau of Supplies and Accounts, Chief Navy Department Washington, D.C. Attn: Code W3	1

<u>Organization</u>	<u>No. of Copies</u>
U.S. Naval Avionics Facility Indianapolis 18, Indiana Attn: Librarian, Code 031.2	1
National Aeronautics & Space Administration Goddard Space Flight Center Greenbelt, Maryland Attn: Chief, Data Systems Div., C.V.L. Smith	1
Federal Aviation Agency Bureau of Research and Development Washington 25, D.C. Attn: RD-375, Mr. Harry Hayman	1
Cornell Aeronautical Laboratory Inc. P. O. Box 235 Buffalo 21, New York Attn: Systems Requirements Dept., A. E. Murray	1
Institute for Space Studies 475 Riverside Drive New York 27, New York Attn: Mr. Albert Arking	1
Stanford University Electronics Laboratory Stanford, California Attn: Prof. N. M. Abramson	1
Lincoln Laboratory Massachusetts Institute of Technology Lexington 73, Massachusetts Attn: Library	1
Electronics Research Laboratory University of California Berkeley 4, California Attn: Director	1
Mr. Gordon Stanley 7685 South Sheridan Ct. Littleton, Colorado Attn: Martin, Denver	1

<u>Organization</u>	<u>No. of Copies</u>
L. Verbeek C.C.R. - Euratom C.E.T.I.S. Casella Postale 1, Ispra, Varese, ITALY	1
A. J. Cote, Jr. Applied Physics Laboratory, JHU The Johns Hopkins University 8621 Georgia Avenue Silver Spring, Maryland	1
Institute for Defense Analysis Communications Research Division Von Neumann Hall Princeton, New Jersey	1
Dr. J. H. Andreae Standard Telecommunication Laboratories Limited London Road, Harlow, Essex ENGLAND	1
Allan Kiron Office of Research & Development U.S. Department of Commerce Patent Office Washington 25, D.C.	1
Air Force Office of Scientific Research Information Research Div. Washington 25, D.C. Attn: R. W. Swanson	1

UNCLASSIFIED

UNCLASSIFIED